

# Leptogenesis beyond the limit of hierarchical heavy neutrino masses

Steve Blanchet and Pasquale Di Bari

*Max-Planck-Institut für Physik*

*(Werner-Heisenberg-Institut)*

*Föhringer Ring 6, 80805 München, Germany*

February 2, 2008

## Abstract

We calculate the baryon asymmetry of the Universe in thermal leptogenesis beyond the usual lightest right-handed (RH) neutrino dominated scenario ( $N_1$ DS) and in particular beyond the hierarchical limit (HL),  $M_1 \ll M_2 \ll M_3$ , for the RH neutrino mass spectrum. After providing some orientation among the large variety of models, we first revisit the central role of the  $N_1$ DS, with new insights on the dynamics of the asymmetry generation and then discuss the main routes departing from it, focusing on models beyond the HL. We study in detail two examples of ‘strong-strong’ wash-out scenarios: one with ‘maximal phase’ and the limit of very large  $M_3$ , studying the effects arising when  $\delta_2 \equiv (M_2 - M_1)/M_1$  is small. We extend analytical methods already applied to the  $N_1$ DS showing, for example, that, in the degenerate limit (DL), the efficiency factors of the RH neutrinos become equal with the single decay parameter replaced by the sum. Both cases disprove the misconception that close RH neutrino masses necessarily lead to a final asymmetry enhancement and to a relaxation of the lower bounds on  $M_1$  and on the initial temperature of the radiation-dominated expansion. We also explain why leptogenesis tends to favor normal hierarchy compared to inverted hierarchy for the left-handed neutrino masses.

# 1 Introduction

With the discovery of neutrino masses in neutrino mixing experiments, leptogenesis [1] has become one of the most attractive explanations of the matter-antimatter asymmetry of the Universe. Indeed, leptogenesis is the direct cosmological consequence of the see-saw mechanism [2], the most elegant way to understand neutrino masses and their lightness compared to all other known massive fermions. Adding to the Standard Model Lagrangian three RH neutrinos with Yukawa coupling matrix  $h$  and Majorana mass matrix  $M$ , a neutrino Dirac mass matrix  $m_D = h v$  is generated, after electroweak symmetry breaking, by the vev  $v$  of the Higgs boson. For  $M \gg m_D$ , the neutrino mass spectrum splits into 3 heavy Majorana states  $N_1, N_2$  and  $N_3$  with masses  $M_1 \leq M_2 \leq M_3$ , which almost coincide with the eigenvalues of  $M$ , and 3 light Majorana states with masses  $m_1 \leq m_2 \leq m_3$  corresponding to the eigenvalues of the neutrino mass matrix given by the see-saw formula,

$$m_\nu = -m_D \frac{1}{M} m_D^T. \quad (1)$$

Neutrino mixing experiments measure two mass-squared differences. In normal (inverted) neutrino schemes, one has

$$m_3^2 - m_2^2 = \Delta m_{\text{atm}}^2 \quad (\Delta m_{\text{sol}}^2), \quad (2)$$

$$m_2^2 - m_1^2 = \Delta m_{\text{sol}}^2 \quad (\Delta m_{\text{atm}}^2 - \Delta m_{\text{sol}}^2). \quad (3)$$

For  $m_1 \gg m_{\text{atm}} \equiv \sqrt{\Delta m_{\text{atm}}^2 + \Delta m_{\text{sol}}^2} \simeq 0.05 \text{ eV}$ , one has a quasi-degenerate spectrum with  $m_1 \simeq m_2 \simeq m_3$ , whereas for  $m_1 \ll m_{\text{sol}} \equiv \sqrt{\Delta m_{\text{sol}}^2} \simeq 0.009 \text{ eV}$  one has a fully hierarchical (normal or inverted) spectrum.

A lepton asymmetry can be generated from the decays of the heavy neutrinos into leptons and Higgs bosons and partly converted into a baryon asymmetry by the sphaleron ( $B-L$  conserving) processes at temperatures higher than about 100 GeV. The asymmetry produced by each  $N_i$  decay is given by the  $CP$  asymmetry parameter  $\varepsilon_i$

$$\varepsilon_i \equiv - \frac{\Gamma_i - \bar{\Gamma}_i}{\Gamma_i + \bar{\Gamma}_i}, \quad (4)$$

where  $\Gamma_i$  is the decay rate into leptons and  $\bar{\Gamma}_i$  the one into anti-leptons. For each  $N_i$  one introduces the decay parameter  $K_i$ , defined as the ratio of the total decay width to the expansion rate at  $T = M_i$ ,

$$K_i \equiv \frac{\tilde{\Gamma}_i}{H(T = M_i)}. \quad (5)$$

This is the key quantity for the thermodynamical description of the decays of heavy particles in the early Universe [3]. In leptogenesis it can be conveniently expressed in

terms of the *effective neutrino mass*  $\tilde{m}_i \equiv (m_D^\dagger m_D)_{ii}/M_i$ , such that  $K_i = \tilde{m}_i/m_\star$ , where  $m_\star \simeq 1.08 \times 10^{-3} \text{ eV}$  is the *equilibrium neutrino mass*. Besides decays, there are other processes, especially inverse decays, that are relevant not only for producing the RH neutrinos but also for washing-out part of the asymmetry produced from decays. The effect of production and wash-out are simultaneously accounted for by the efficiency factors  $\kappa_i$  associated to the production of the asymmetry from each  $N_i$ , such that the final  $B-L$  asymmetry can be expressed as the sum of three contributions

$$N_{B-L}^f = \sum_i \varepsilon_i \kappa_i^f. \quad (6)$$

The baryon-to-photon number ratio at the recombination time can then be calculated as

$$\eta_B = a_{\text{sph}} \frac{N_{B-L}^f}{N_\gamma^{\text{rec}}} \simeq 0.96 \times 10^{-2} \sum_i \varepsilon_i \kappa_i^f, \quad (7)$$

where  $a_{\text{sph}} \simeq 1/3$  is the sphaleron conversion coefficient. Here we assume a standard thermal history and indicate with  $N_X$  any particle number or asymmetry  $X$  calculated in a portion of comoving volume containing one heavy neutrino in ultra-relativistic thermal equilibrium, so that  $N_\gamma^{\text{rec}} \simeq 37$ . The efficiency factors  $\kappa_i^f \rightarrow 1$  in the limit of an initial ultra-relativistic thermal  $N_i$  abundance and null wash-out.

A great simplification occurs in the HL<sup>1</sup> ( $M_1 \ll M_2 \ll M_3$ ). In this case one typically (but not necessarily!) obtains what can be called the  $N_1$ DS, where both the wash-out from the two heavier RH neutrinos and the asymmetry produced by their decays can be neglected and the expression (6) reduces to  $(N_{B-L}^f)_{\text{HL}} \simeq \kappa_1^f \varepsilon_1$ . The HL is quite a natural assumption for hierarchical light neutrinos.

The effective neutrino mass  $\tilde{m}_1$  can be expressed as a linear combination of the light neutrino masses with positive coefficients whose sum cannot be smaller than unity. For this reason, the experimental findings  $m_{\text{sol}}, m_{\text{atm}} \gg m_\star$  typically force  $K_1$  to lie in the range  $\mathcal{O}(K_{\text{sol}} \simeq 9) \lesssim K_1 \lesssim \mathcal{O}(K_{\text{atm}} \simeq 50)$ , where  $K_{\text{sol}} \equiv m_{\text{sol}}/m_\star$  and  $K_{\text{atm}} \equiv m_{\text{atm}}/m_\star$ , i.e. in the strong wash-out regime ( $K_1 \gg 1$ ), while the weak wash-out regime ( $K_1 \lesssim 1$ ) is possible for a particular class of neutrino mass models.

The efficiency factor  $\kappa_1^f$  is approximately given by the *number of  $N_1$  that decay out-of-equilibrium*. In the strong wash-out regime this is unambiguously specified by the thermal equilibrium abundance at the time when the inverse decays get frozen, at a well-defined temperature  $T_B \ll M_1$  when the  $N_1$ 's are non-relativistic [4]. One has then to require

---

<sup>1</sup>Throughout the paper we make use of the following acronyms: HL=hierarchical limit, DL=degenerate limit,  $N_i$ DS= $N_i$ -dominated scenario.

that the initial temperature of the Universe is larger than  $\sim T_B$ , the key assumption for thermal leptogenesis. Therefore, in the strong wash-out regime only a small fraction of  $N_1$ , compared to an initial ultra-relativistic thermal abundance, decays out-of-equilibrium. This results in small values for  $\kappa_1^f \sim 10^{-3} \div 10^{-2}$ , but still large enough to allow for successful leptogenesis in quite a large region of parameter space. On the other hand, the positive by-product of the strong wash-out regime is that the final asymmetry does not depend on the initial conditions.

A second important simplification occurring in the HL is that the  $CP$  asymmetry  $\varepsilon_1$ , like  $\kappa_1^f$ , depends only on a limited set of see-saw parameters and, quite remarkably, it turns out that there is an upper bound on  $\varepsilon_1$  proportional to  $M_1$  [5, 6]. In the strong wash-out regime for  $K_1 \gtrsim 5$ , this gives rise to a lower bound on the lightest RH neutrino mass  $M_1 \gtrsim 5 \times 10^9 \text{ GeV}$  [7, 6, 8], also implying a lower bound on the initial temperature of the radiation-dominated expansion  $T_{\text{in}} \gtrsim 2 \times 10^9 \text{ GeV}$  [4, 8]<sup>2</sup>, identifiable with the reheating temperature within inflation [11]. In the  $N_1\text{DS}$ , for quasi-degenerate neutrinos, the combined effect [12] of the additional wash-out from  $\Delta L = 2$  processes, which depends on the combination  $M_1 \sum m_i^2$ , together with a  $CP$  asymmetry suppression [6, 13], gives rise to a stringent upper bound on the absolute neutrino mass scale  $m_1 \leq 0.1 \text{ eV}$  [12, 13, 9, 4].

A weak version of the  $N_1\text{DS}$ , for  $K_1 \lesssim 1$ , encounters two serious difficulties. The first is the strong dependence on the initial conditions, preventing the model from being self-contained. The second is that, for  $\tilde{m}_1 \rightarrow m_1$ , the  $CP$  asymmetry vanishes and  $\tilde{m}_1$  has to be fine-tuned to have successful leptogenesis. A more appealing possibility is then represented by the  $N_2\text{DS}$  [8], where the asymmetry is mostly generated from the decays of  $N_2$ , circumventing both problems.

Another key motivation to study models beyond the HL is to allow RH neutrino masses to be arbitrarily close. This possibility has been considered in many works [14, 15, 16]. In [8] an analytical condition for the validity of a calculation of the efficiency factor  $\kappa_1^f$  in the HL was found. It was noticed that one may neglect the effect of the two heavier neutrinos if  $\delta_2 \equiv (M_2 - M_1)/M_1 \gtrsim 1.5 \div 5$ , the exact value depending on  $K_1$  and  $K_2$ . The validity of the HL in the calculation of the  $CP$  asymmetry is more involved but a similar condition holds in most cases. There is one possibility, discussed in the case C of Section 4, where the condition on  $\delta_2$  to recover the HL becomes more stringent. For quasi-degenerate neutrinos this also provides a way to evade the upper bound on neutrino masses [17].

In this paper we perform a general calculation of the final asymmetry beyond the

---

<sup>2</sup>In the MSSM these values become respectively  $M_1 \gtrsim 2 \times 10^9 \text{ GeV}$  and  $T_{\text{in}} \gtrsim 10^9 \text{ GeV}$  [9, 10, 8].

HL, extending useful analytical methods described in [4] within the HL. A discussion of flavor effects [18, 19, 20] is deferred to a forthcoming paper, since they are somewhat complementary to the issues addressed here. However, it is worthwhile to mention that, accounting for these effects, the upper bound on the absolute neutrino mass scale does not hold [20].

The main difficulty of such a general calculation is the great model dependence. Therefore, in Section 2 our first step is a description of a general way to parameterize and classify models. In Section 3 we revisit the  $N_1$ DS, providing several new interesting analytical insights on the dynamics of the asymmetry generation. In Section 4 we describe the main routes to go beyond the  $N_1$ DS, including models beyond the HL that we study in detail in Section 5. Here we focus on *strong-strong* wash-out scenarios, where both  $K_1$  and  $K_2 \gtrsim 5$  and  $M_3 \gg M_1, M_2$ , but with arbitrary  $M_1$  and  $M_2$ . We show how the production and the wash-out from each RH neutrino interfere with each other, calculating the efficiency factors and giving exact conditions for the HL to be recovered. Then we calculate the lower bounds on  $M_1$  and  $T_{\text{in}}$  first in a model where the asymmetry is maximal in the HL, but insensitive to a  $CP$  asymmetry enhancement beyond the HL, then in a model that received recently great attention, where  $M_3 \gg 10^{14} \text{ GeV} \gg M_2, M_1$  [21]. We also explain why leptogenesis favors normal hierarchy over an inverted one. We summarize our conclusions in Section 6.

## 2 Getting oriented among leptogenesis scenarios

Let us describe how to calculate the final asymmetry for a general RH neutrino spectrum. In a one-flavor approximation, the set of kinetic equations can be written as [22, 14, 15]

$$\frac{dN_{N_i}}{dz} = -(D_i + S_i) (N_{N_i} - N_{N_i}^{\text{eq}}), \quad i = 1, 2, 3 \quad (8)$$

$$\frac{dN_{B-L}}{dz} = \sum_{i=1}^3 \varepsilon_i D_i (N_{N_i} - N_{N_i}^{\text{eq}}) - N_{B-L} W, \quad (9)$$

where  $z \equiv M_1/T$ . Defining  $x_i \equiv M_i^2/M_1^2$  and  $z_i \equiv z \sqrt{x_i}$ , the *decay factors* are given by

$$D_i \equiv \frac{\Gamma_{D,i}}{H z} = K_i x_i z \left\langle \frac{1}{\gamma_i} \right\rangle, \quad (10)$$

where  $H$  is the expansion rate. The total decay rates,  $\Gamma_{D,i} \equiv \Gamma_i + \bar{\Gamma}_i$ , are the product of the decay widths times the thermally averaged dilation factors  $\langle 1/\gamma_i \rangle$ , given by the ratio  $\mathcal{K}_1(z_i)/\mathcal{K}_2(z_i)$  of the modified Bessel functions. The equilibrium abundance and its rate

are also expressed through the modified Bessel functions,

$$N_{N_i}^{\text{eq}}(z_i) = \frac{1}{2} z_i^2 \mathcal{K}_2(z_i) \quad , \quad \frac{dN_{N_i}^{\text{eq}}}{dz_i} = -\frac{1}{2} z_i^2 \mathcal{K}_1(z_i) . \quad (11)$$

The RH neutrinos can be produced by inverse decays and scatterings. Nevertheless, in the relevant strong wash-out regime, the inverse decays alone are already sufficient to make the RH neutrino abundance reach its thermal equilibrium value prior to their decays. Therefore, the details of the RH neutrino production do not affect the final asymmetry and theoretical uncertainties are consequently greatly reduced. This is one of the nice features of the strong wash-out regime on which we will focus and for this reason the scattering terms  $S_i$  will play no role. The wash-out factor  $W$  can be written as the sum of two contributions [7],

$$W = \sum_i W_i(K_i) + \Delta W(M_1 \bar{m}^2) . \quad (12)$$

The second term arises from the non-resonant  $\Delta L = 2$  processes and gives typically a non-negligible contribution only in the non relativistic limit for  $z \gg 1$  [22, 7, 4]. For hierarchical light neutrinos it can be safely neglected for reasonable values  $M_1 \ll 10^{14}$  GeV. In the strong wash-out regime, the first term is dominated by inverse decays [9, 4], where the resonant  $\Delta L = 2$  contribution has to be properly subtracted [23, 9], so that <sup>3</sup>

$$W_i(z) \simeq W_i^{\text{ID}}(z) = \frac{1}{4} K_i \sqrt{x_i} \mathcal{K}_1(z_i) z_i^3 . \quad (13)$$

Let us indicate with  $N_{B-L}^{\text{in}}$  a possible pre-existing asymmetry at  $T_{\text{in}}$ . The final asymmetry can then be written in an integral form [3, 4] ,

$$N_{B-L}^{\text{f}} = N_{B-L}^{\text{in}} e^{-\sum_i \int dz' W_i(z')} + \sum_i \varepsilon_i \kappa_i^{\text{f}} , \quad (14)$$

with the efficiency factors  $\kappa_i^{\text{f}}$  given by

$$\kappa_i^{\text{f}} = - \int_{z_{\text{in}}}^{\infty} dz' \frac{dN_{N_i}}{dz'} e^{-\sum_i \int_{z'}^z dz'' W_i(z'')} , \quad (15)$$

where we defined  $z_{\text{in}} \equiv M_1/T_{\text{in}}$ . Notice that, in general, each efficiency factor depends on all decay parameters, i.e.  $\kappa_i^{\text{f}} = \kappa_i^{\text{f}}(K_1, K_2, K_3)$ .

If the mass differences satisfy the condition for the applicability of perturbation theory,  $|M_j - M_i|/M_i \gg \max[(h^\dagger h)_{ij}]/(16\pi^2)$  with  $j \neq i$  [24], then a perturbative calculation from the interference of tree level with one loop self-energy and vertex diagrams gives [25]

$$\varepsilon_i = \frac{3}{16\pi} \sum_{j \neq i} \frac{\text{Im} [(h^\dagger h)_{ij}^2]}{(h^\dagger h)_{ii}} \frac{\xi(x_j/x_i)}{\sqrt{x_j/x_i}} \quad (16)$$

---

<sup>3</sup>In the following we will imply this subtraction when referring to the ‘wash-out from inverse decays’.




Figure 1: The function  $\xi(x)$  defined in the Eq. (17).

where the function  $\xi(x)$ , shown in Fig. 1, is defined as [13]

$$\xi(x) = \frac{2}{3} x \left[ (1+x) \ln \left( \frac{1+x}{x} \right) - \frac{2-x}{1-x} \right]. \quad (17)$$

Working in a basis where both the Majorana mass and the light neutrino mass matrix are diagonal,  $D_M \equiv \text{diag}(M_1, M_2, M_3)$  and  $D_m \equiv \text{diag}(m_1, m_2, m_3) = -U^\dagger m_\nu U^*$ , from the see-saw formula one can obtain a useful parametrization of the Dirac mass matrix in terms of the orthogonal complex matrix  $\Omega$  [26]<sup>4</sup>,

$$m_D = U \sqrt{D_m} \Omega \sqrt{D_M}. \quad (18)$$

The unitary matrix  $U$  can be identified with the PMNS matrix in a basis where the charged lepton mass matrix is also diagonal. The following parametrization of the  $\Omega$  matrix proves to be particularly useful in leptogenesis [8]

$$\Omega(\Omega_{21}, \Omega_{31}, \Omega_{22}) = \begin{pmatrix} \sqrt{1 - \Omega_{21}^2 - \Omega_{31}^2} & \Omega_{12} & \pm \sqrt{\Omega_{21}^2 + \Omega_{31}^2 - \Omega_{12}^2} \\ \Omega_{21} & \Omega_{22} & -\sqrt{1 - \Omega_{22}^2 - \Omega_{21}^2} \\ \Omega_{31} & \sqrt{1 - \Omega_{22}^2 - \Omega_{12}^2} & \sqrt{\Omega_{22}^2 + \Omega_{12}^2 - \Omega_{31}^2} \end{pmatrix}, \quad (19)$$

where  $\Omega_{12}$  can be expressed as a function of  $(\Omega_{21}, \Omega_{31}, \Omega_{22})$  imposing, for example,  $\sum_j \Omega_{j1} \Omega_{j2} = 0$ . From this general form one can obtain, for particular choices of the parameters, three elementary complex rotations that exhibit peculiar properties. In terms

<sup>4</sup>Compared to the  $R$  matrix in [26], one has the simple relation  $\Omega = R^\dagger$ .

of these rotations an alternative parametrization of the  $\Omega$  matrix is

$$\Omega(\omega_{21}, \omega_{31}, \omega_{22}) = R_{12}(\omega_{21}) R_{13}(\omega_{31}) R_{23}(\omega_{22}) , \quad (20)$$

where

$$R_{12} = \begin{pmatrix} \sqrt{1 - \omega_{21}^2} & -\omega_{21} & 0 \\ \omega_{21} & \sqrt{1 - \omega_{21}^2} & 0 \\ 0 & 0 & 1 \end{pmatrix} , \quad R_{13} = \begin{pmatrix} \sqrt{1 - \omega_{31}^2} & 0 & -\omega_{31} \\ 0 & 1 & 0 \\ \omega_{31} & 0 & \sqrt{1 - \omega_{31}^2} \end{pmatrix} , \quad R_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega_{22} & -\sqrt{1 - \omega_{22}^2} \\ 0 & \sqrt{1 - \omega_{22}^2} & \omega_{22} \end{pmatrix} . \quad (21)$$

This parametrization for an orthogonal complex matrix corresponds to the transposed form of the CKM matrix in the quark sector or of the PMNS unitary matrix in neutrino mixing, with the difference that here one has complex rotations instead of real ones. There are straightforward relations between the parameters  $\Omega_{ij}$  and the  $\omega_{ij}$ 's:

$$\Omega_{21} = \omega_{21} \sqrt{1 - \omega_{31}^2} , \quad \Omega_{31} = \omega_{31} , \quad \Omega_{22} = \omega_{22} \sqrt{1 - \omega_{21}^2} - \omega_{21} \omega_{31} \sqrt{1 - \omega_{22}^2} . \quad (22)$$

The two parameterizations are interchangeable and it can be more convenient to use one or the other depending on the context. The parametrization Eq. (20) is particularly useful to understand the general structure of different models occurring in thermal leptogenesis.

- For  $\Omega = R_{13}$  one has  $\varepsilon_2 = 0$ , while  $\varepsilon_1$  is maximal if [8]  $m_3 \text{Re}(\omega_{31}^2)/|\omega_{31}^2| = m_1 [1 - \text{Re}(\omega_{31}^2)]/|1 - \omega_{31}^2|$ . In the HL ( $M_2 \gg M_1$ ) one obtains the  $N_1$ DS.
- For  $\Omega = R_{23}$  one has  $\varepsilon_1 = 0$ , while  $\varepsilon_2$  is maximal if  $m_3 \text{Re}(\omega_{22}^2)/|\omega_{22}^2| = m_2 [1 - \text{Re}(\omega_{22}^2)]/|1 - \omega_{22}^2|$ . At the same time, one has  $\tilde{m}_1 = m_1$ , so that the wash-out from  $N_1$  can be neglected if  $m_1$  is small enough. Therefore, in the HL and for hierarchical light neutrinos, one obtains the  $N_2$ DS [8], as discussed in Section 4.
- If  $\Omega = R_{12}$ , then  $\varepsilon_1$  undergoes a phase suppression compared to its maximal value but  $|\varepsilon_2| \propto (M_1/M_2) |\varepsilon_1|$ . This implies that in the HL one again recovers the  $N_1$ DS [8]. On the other hand if  $M_1 \simeq M_2$  both  $CP$  asymmetries can play a role.

Notice that  $N_2$ DS requires a more special  $\Omega$  form than the  $N_1$ DS, but on the other hand there is no lower bound on  $M_1$  as in the  $N_1$ DS. Therefore, it represents an interesting alternative [8]. On the other hand, there cannot be a  $N_3$ DS with only 3 RH neutrinos. The reason is that  $\varepsilon_3 \rightarrow 0$  in the HL for any  $\Omega$ . This can be understood more generally if one observes that the  $CP$  asymmetry of a decaying particle vanishes in the limit where all particles in the propagators are massless. This also explains why it is more special to have  $|\varepsilon_2| \gg |\varepsilon_1|$  than the opposite: in the first case one must necessarily have  $N_3$  running in the propagator in order to have  $\varepsilon_2 \neq 0$  and this happens for  $\Omega = R_{23}$ , while in order to have  $\varepsilon_1 \neq 0$  one can have either  $N_2$  for  $\Omega = R_{12}$ , or  $N_3$  for  $\Omega = R_{13}$ , or both.



### 3 Revisiting the $N_1$ DS

Let us now revisit shortly some of the results holding in the  $N_1$ DS, with some new interesting insights on the dynamics of the asymmetry generation. This is also necessary to introduce quantities and notations that will be used or extended in Section 5 to go beyond the HL.

The general expression for the final asymmetry Eq. (7) reduces to  $\eta_B \simeq 10^{-2} \varepsilon_1 \kappa_1^f$ , where  $\kappa_1^f$  can be calculated solving a system of just two kinetic equations [14, 7, 9], obtained from the general set (cf. (8) and (9)) neglecting the asymmetry generation and the wash-out terms from the two heavier RH neutrinos.

For  $M_1 \ll 10^{14} \text{ GeV} (m_{\text{atm}}^2 / \sum m_i^2)$ , the term  $\Delta W(z)$  in the wash-out (cf. (12)) is negligible and the solutions depend just on  $K_1$ , since this is the only parameter in the equations. They can be worked out in an integral form [3] and for the  $B - L$  asymmetry one obtains a special case of the more general Eq. (14),

$$N_{B-L}(z; \bar{z}) = \bar{N}_{B-L} e^{-\int_{\bar{z}}^z dz' W_1(z')} + \varepsilon_1 \kappa_1(z; \bar{z}), \quad (23)$$

where now a possible asymmetry produced from the two heavier RH neutrinos and frozen at  $\bar{z} \geq z_{\text{in}}$  is included in  $\bar{N}_{B-L}$ . The efficiency factor  $\kappa_1(z; \bar{z})$  can be expressed through a Laplace integral,

$$\kappa_1(z; \bar{z}) = - \int_{\bar{z}}^z dz' \frac{dN_{N_1}}{dz'} e^{-\int_{z'}^{\infty} dz'' W_1(z'')} = \int_{\bar{z}}^z dz' e^{-\psi(z', z)}. \quad (24)$$

In the strong wash-out regime, using the approximation  $dN_{N_1}/dz' \simeq dN_{N_1}^{\text{eq}}/dz'$  and  $W_1(z') \simeq W_1^{\text{ID}}(z')$  (cf. (13)), one finds that for  $z \rightarrow \infty$  the final value is given by [4]

$$\kappa_1^f(K_1) \simeq \kappa(K_1) \equiv \frac{2}{K_1 z_B(K_1)} \left( 1 - e^{-\frac{K_1 z_B(K_1)}{2}} \right), \quad (25)$$

if  $\bar{z} \lesssim z_B - 2$ . The value  $z' = z_B(K_1)$  is where the quantity  $\psi(z', \infty)$  has a minimum and the integral in the Eq. (24) receives a dominant contribution from a restricted  $z'$ -interval centered around it. In the strong wash-out regime it is well reproduced by

$$z_B(K_1) \simeq 2 + 4 K_1^{0.13} e^{-\frac{2.5}{K_1}}. \quad (26)$$

Figures 2 and 3 show, for an initial thermal abundance, the dynamics of the asymmetry generation, comparing one example of weak wash-out with one example of strong wash-out<sup>5</sup>. In the top panels we show the function  $d\kappa_1/dz' \equiv e^{-\psi(z', z)}$ , defined for  $z' \leq z$ , for different values of  $z$ . The difference between the two cases is striking.




Figure 2: Dynamics in the weak wash-out regime for initial thermal abundance ( $N_{N_1}^{\text{in}} = 1$ ). Top panel: rates. Bottom panel: efficiency factor  $\kappa_1$  and  $N_1$ -abundance. The maximum of the asymmetry production rate occurs at  $z' \simeq z_{\max}^{\text{weak}} = 1/\sqrt{K_1} + 15/8 \simeq 12$ .

In the weak wash-out regime, each decay contributes to the final asymmetry for any value of  $z'$  when the asymmetry is produced. In the strong wash-out regime, all asymmetry produced at  $z' \lesssim z_B - 2$  is efficiently washed-out by inverse decays, such that only decays occurring at  $z' \sim z_B$  give a contribution to the final asymmetry.

<sup>5</sup>Corresponding animations can be found at <http://wwwth.mppmu.mpg.de/members/dibari>.




Figure 3: Dynamics in the strong wash-out regime. Top panel: rates. Bottom panel: efficiency factor  $\kappa_1$  and  $N_1$ -abundance. The maximum of the final asymmetry production rate occurs at  $z' \simeq z_B$ .

The general expression Eq. (16) for  $\varepsilon_1$  can be re-cast through the  $\Omega$  matrix as [8]

$$\varepsilon_1 = \xi(x_2) \varepsilon_1^{\text{HL}}(m_1, M_1, \Omega_{21}, \Omega_{31}) + [\xi(x_3) - \xi(x_2)] \Delta \varepsilon_1(m_1, M_1, \Omega_{21}, \Omega_{31}, \Omega_{22}). \quad (27)$$

In the HL, for  $x_3, x_2 \gg 1$ , one has  $\xi(x_2) \simeq \xi(x_3) \simeq 1$  and then, from Eq. (27),

$\varepsilon_1 = \varepsilon_1^{\text{HL}}(m_1, M_1, \Omega_{21}, \Omega_{31})$ . Therefore, in the HL the dependence on four of the see-saw parameters, namely  $M_2, M_3, \Omega_{22}$ , cancels out and one is left only with six parameters. Notice moreover that  $\kappa_1^f = \kappa_1^f(m_1, M_1, K_1)$ , where  $K_1 = K_1(m_1, \Omega_{21}, \Omega_{31})$ , and thus the final asymmetry depends on the same six parameters too.

The HL for  $\varepsilon_1$  can be written as [6]  $\varepsilon_1^{\text{HL}}(m_1, M_1, \Omega_{21}, \Omega_{31}) \equiv \bar{\varepsilon}(M_1) \beta(m_1, \Omega_{21}, \Omega_{31})$ , where, writing  $\Omega_{ij}^2 \equiv X_{ij} - i Y_{ij} \equiv \rho_{ij} e^{i \varphi_{ij}}$ , with  $\rho_{ij} \equiv |\Omega_{ij}^2| \geq 0$ , one has <sup>6</sup>

$$\bar{\varepsilon}(M_1) \equiv \frac{3}{16\pi} \frac{M_1 m_{\text{atm}}}{v^2} \quad \text{and} \quad \beta(m_1, \Omega_{21}, \Omega_{31}) \equiv \frac{\sum_j m_j^2 Y_{j1}}{m_{\text{atm}} \sum_j m_j \rho_{j1}}. \quad (28)$$

It is interesting that  $\beta(m_1, \Omega_{21}, \Omega_{31}) \leq 1$ , so that in the HL one has the upper bound  $|\varepsilon_1^{\text{HL}}| \leq \bar{\varepsilon}(M_1)$  [5, 6]. More precisely one can define an *effective phase*  $\delta_L^{(1)}$  by

$$\beta(m_1, \Omega_{21}, \Omega_{31}) = \beta_{\text{max}}(m_1, \tilde{m}_1) \sin \delta_L^{(1)}(m_1, \Omega_{21}, \Omega_{31}), \quad (29)$$

such that the upper bound [6, 13]

$$\beta_{\text{max}}(m_1, \tilde{m}_1) = \frac{m_{\text{atm}}}{m_1 + m_3} f(m_1, \tilde{m}_1) \leq 1, \quad (30)$$

corresponds to  $\sin \delta_L^{(1)} = 1$  and it is obtained by maximizing over the  $\Omega$ -parameters for fixed  $\tilde{m}_1$ . The function  $f(m_1, \tilde{m}_1)$  is [8]

$$f(m_1, \tilde{m}_1) = \frac{m_1 + m_3}{\tilde{m}_1} Y_{\text{max}}(m_1, \tilde{m}_1), \quad (31)$$

where  $Y_{\text{max}}(m_1, \tilde{m}_1)$  is the maximum of  $Y_{31}$  for  $\Omega_{21} = 0$ . For hierarchical light neutrinos,  $m_1 \lesssim 0.2 m_{\text{atm}}$ , an approximate explicit expression is [13]

$$f(m_1, \tilde{m}_1) = \frac{m_3 - m_1 \sqrt{1 + \frac{m_3^2 - m_1^2}{\tilde{m}_1^2}}}{m_3 - m_1}, \quad (32)$$

which further simplifies to  $f(m_1, \tilde{m}_1) = 1 - m_1/\tilde{m}_1$  for  $m_1 \ll 0.1 m_{\text{atm}}$ . Conversely, in the quasi-degenerate limit, one has  $f(m_1, \tilde{m}_1) = \sqrt{1 - (m_1/\tilde{m}_1)^2}$  [17]. One can then conclude that the maximum of the  $CP$  asymmetry is reached in the limit  $m_1 \rightarrow 0$ , when  $f(m_1, \tilde{m}_1) = 1$  and, since this is true also for  $\kappa_1^f$ , it applies also to the final asymmetry  $N_{B-L}^f \simeq \varepsilon_1 \kappa_1^f$ . For  $\Omega_{21} = 0$  and  $Y_{31} = Y_{\text{max}}(m_1, \tilde{m}_1)$ , the phase is maximal, while for a generic choice of  $\Omega$ , the  $CP$  asymmetry undergoes a phase suppression

$$\sin \delta_L^{(1)}(m_1, \Omega_{21}, \Omega_{31}) = \frac{m_1 + m_3}{\tilde{m}_1 f(m_1, \tilde{m}_1)} (Y_{31} + \sigma^2 Y_{21}) = \frac{(Y_{31} + \sigma^2 Y_{21})}{Y_{\text{max}}(m_1, \tilde{m}_1)}, \quad (33)$$

---

<sup>6</sup>In [8] a minus sign in the expression for  $\beta(m_1, \Omega_{21}, \Omega_{31})$  is missing. This does not affect any of the results but the quantities  $Y_{i1}$  should be defined as  $Y_{i1} \equiv -\text{Im}(\Omega_{j1}^2)$  as here.

where  $\sigma \equiv \sqrt{m_2^2 - m_1^2}/m_{\text{atm}}$ . One can see that  $\sin \delta_L^{(1)} = 1$  for  $Y_{21} = 0$  and  $Y_{31} = Y_{\text{max}}$ . It is instructive to calculate  $\sin \delta_L^{(1)}$  for each of the three elementary complex rotations that can be used to parameterize  $\Omega$  (cf. 20)):

- For  $\Omega = R_{13}$ , one has  $\sin \delta_L^{(1)} = Y_{31}/Y_{\text{max}}$  and the phase is maximal if  $Y_{31} = Y_{\text{max}} = \tilde{m}_1/m_{\text{atm}}$ ; notice that there is no difference between normal and inverted hierarchy.
- For  $\Omega = R_{12}$ , one has  $\sin \delta_L^{(1)} = \sigma^2 Y_{21}/Y_{\text{max}} \leq \sigma$ , larger for inverted hierarchy compared to the normal one; however since  $\kappa_1^f \propto K_1^{-1.2}$  [10] and since  $K_1 = K_{\text{sol}} |\Omega_{21}^2|$  for normal hierarchy and  $K_1 \simeq K_{\text{atm}} |\Omega_{21}^2|$  for inverted hierarchy, the final asymmetry is slightly higher for normal hierarchy compared to the inverted one for fixed  $|\Omega_{21}^2|$  [8].
- For  $\Omega = R_{23}$ , one has  $\sin \delta_L^{(1)} = \varepsilon_1 = 0$ ; one can check that  $\varepsilon_1 = 0$  applies independently of  $M_2$  and  $M_3$  and therefore not only in the HL. Notice that the conclusions in the previous two cases are still valid if one multiplies  $R_{13}$  or  $R_{12}$  with  $R_{23}$  respectively, since it does not affect  $\sin \delta_L^{(1)}$ .

Interesting constraints follow if one imposes that the asymmetry produced from leptogenesis explains the measured value from WMAP plus SLOAN combined determination [7, 27],

$$\eta_B(m_1, M_1, \Omega_{21}, \Omega_{31}) = \eta_B^{\text{CMB}} = (6.3 \pm 0.3) \times 10^{-10}. \quad (34)$$

If  $M_1 \ll 10^{14} \text{ GeV}$  ( $m_{\text{atm}}^2/\sum_i m_i^2$ ), then

$$M_1 \simeq \frac{N_\gamma^{\text{rec}}}{a_{\text{sph}}} \frac{16 \pi v^2}{3} \frac{\eta_B^{\text{CMB}}}{m_{\text{atm}}} \left[ \kappa(K_1) \beta_{\text{max}}(m_1, K_1) \sin \delta_L^{(1)}(\Omega_{21}, \Omega_{31}) \right]^{-1} \quad (35)$$

$$\gtrsim (M_1^{\text{min}})_{\text{HL}} \equiv \frac{4.2 \times 10^8 \text{ GeV}}{\kappa(K_1) \beta_{\text{max}}(m_1, K_1) \sin \delta_L^{(1)}(\Omega_{21}, \Omega_{31})} \quad (\text{at } 3\sigma \text{ C.L.}) \quad (36)$$

This expression is quite general and shows the effect of the phase suppression [8] and of a higher absolute neutrino mass scale [10] in making the lower bound more restrictive. In Fig. 4 we show  $M_1^{\text{min}}$  (thick solid line) for fully hierarchical light neutrinos ( $m_1 = 0$ ) and maximal phase ( $\sin \delta_L = 1$ ). For  $K_1 \gtrsim 5$  one obtains the lowest value independent of the initial conditions [8],

$$M_1 \gtrsim 5 \times 10^9 \text{ GeV}. \quad (37)$$

The lower bound on  $M_1$  also translates into a lower bound on  $T_{\text{in}}$ ,

$$T_{\text{in}} \geq (T_{\text{in}}^{\text{min}})_{\text{HL}} \simeq \frac{(M_1^{\text{min}})_{\text{HL}}}{z_B(K_1) - 2} \gtrsim 2 \times 10^9 \text{ GeV} \quad (K_1 \gtrsim 5). \quad (38)$$




Figure 4: Lower bounds on  $M_1$  and  $T_{\text{in}}$  vs.  $K_1$  (cf. (36) and (38)). Thick lines: case of maximal phase ( $\sin \delta_L^{(1)} = 1$ ). Thin lines: case of very heavy  $N_3$ .

A plot of this lower bound is shown in Fig. 4 (thick dashed line). The relation between  $M_1^{\text{min}}$  and  $T_{\text{in}}^{\text{min}}$  can be understood from the top panel of Fig. 3, showing that the final asymmetry is the result of the decays occurring just around  $z_B$ , when inverse decays switch off, whereas all asymmetry produced before is efficiently washed out.

## 4 Beyond the $N_1\text{DS}$

There are three ways to go beyond the  $N_1\text{DS}$ . We assume fully hierarchical light neutrinos.

### A) $N_2\text{DS}$

For  $\Omega = R_{23}$ , a nice coincidence is realized: the  $CP$  asymmetry  $\varepsilon_1 = 0$  while  $\varepsilon_2$  can be maximal and, at the same time, the wash-out from the  $N_1$  inverse decays vanishes for  $m_1 \lesssim 10^{-3} \text{ eV}$ . In this way the final asymmetry can and has to be explained in terms of  $N_2$  decays. A nice feature is that the lower bound on  $M_1$  does not hold any more, being replaced by a lower bound on  $M_2$  that, however, still implies a lower bound on  $T_{\text{in}}$  [8]. If one switches on some small  $R_{12}$  and  $R_{13}$  complex rotations, then the lower bounds on  $M_2$  and  $T_{\text{in}}$  become necessarily more stringent. Therefore, there is a border beyond which this scenario is not viable and one is forced to go back to the usual  $N_1\text{DS}$  for

successful leptogenesis. Account of flavor effects contribute to enlarge the domain where the  $N_2$ DS works, since the wash-out from the lightest RH neutrino is diminished [28]. Notice moreover that, like in the  $N_1$ DS, the lower bounds are more stringent for inverted hierarchy than for normal hierarchy, since in the first case  $K_2 \gtrsim K_{\text{atm}}$ , while in the second case  $K_2 \geq K_{\text{sol}}$ .

## B) Beyond the HL

If the RH neutrino masses are sufficiently close then the  $B - L$  asymmetry and the wash-out from the two heavier RH neutrinos have also to be taken into account and the general expression (6) for  $N_{B-L}^f$  has to be used; at the same time, the general expression Eq. (27) for  $\varepsilon_1$  has also to be used. Here the first, typically dominant, term can be enhanced by a factor  $\xi(x_2)$  [13] while the second term can be calculated using [8]

$$\Delta\varepsilon_1 \equiv \frac{3}{16\pi} \frac{\text{Im}[(h^\dagger h)_{13}^2]}{(h^\dagger h)_{11}} \frac{1}{\sqrt{x_3}} = \bar{\varepsilon}(M_1) \frac{\text{Im}[\sum_h m_h \Omega_{h1}^* \Omega_{h3}]^2}{m_{\text{atm}} \tilde{m}_1}. \quad (39)$$

One can see that  $\Delta\varepsilon_1$  vanishes for  $\Omega = R_{12}$ . It also vanishes in another interesting situation, the *strong-strong wash-out scenario* for very large  $M_3$ , that will be discussed in detail in the next Section. In these two interesting cases, the enhancement  $\xi(x_2)$  is the only effect on  $\varepsilon_1$  and the HL is recovered, within a 10% precision, for  $\delta_2 \gtrsim 5$  [13, 8]. These two cases have also been recently studied in [16] for  $\delta_2 \ll 1$ , within the context of radiative leptogenesis.

It is interesting that even the enhancement  $\xi(x_2)$  can be absent in a particular case. This happens if one considers fully hierarchical light neutrinos and  $\Omega = R_{13}$ . It is easy to calculate that in this case one has  $\Delta\varepsilon_1 = \bar{\varepsilon}(M_1) \sin \delta_L^{(1)}$ . Plugging this term into the general expression (27) for  $\varepsilon_1$ , one has an exact cancellation of the terms proportional to  $\xi(x_2)$  and in the end

$$\varepsilon_1|_{\Omega=R_{13}} = \xi(x_3) \bar{\varepsilon}(M_1) \sin \delta_L^{(1)}. \quad (40)$$

For  $x_3 \gg 1$ , such that  $\xi(x_3) \simeq 1$ , there is no enhancement of the usual  $CP$  asymmetry when  $M_2 \rightarrow M_1$ . This example disproves the misconception that degenerate RH neutrino masses unavoidably lead to  $CP$  asymmetry enhancement. Moreover, it shows that the usual most stringent lower bounds on  $M_1$  and  $T_{\text{in}}$ , for  $\sin \delta_L^{(1)} = 1$ , continue to be valid even beyond the HL. However, they can be evaded in other models, which means that, in general, changing the RH neutrino mass spectrum, the  $\Omega$  matrix that maximizes the asymmetry changes too.

As a last exercise, one can check that  $\Delta\varepsilon_1$ , like  $\varepsilon_1^{\text{HL}}$ , vanishes for  $\Omega = R_{23}$ , confirming that the  $N_2$ DS is the only possibility for this particular choice.

### C) Large $|\Omega_{22}|$

There is a situation when the term  $[\xi(x_3) - \xi(x_2)] \Delta \varepsilon_1$  becomes dominant over  $\xi(x_2) \varepsilon_1^{\text{HL}}$  and one has to require a stronger hierarchy to recover the HL [17]. This term is maximized over  $x_3$  for  $x_3 \gg 1$ , implying  $\xi(x_3) \simeq 1$ . Moreover, if for definiteness one imposes  $\Omega_{21} = X_{31} = 0$ , so that  $\sin \delta_L^{(1)} = 1$ , then [8]

$$\xi_{\varepsilon_1} \equiv \frac{\varepsilon_1}{\bar{\varepsilon}(M_1)} = \xi(x_2) + [1 - \xi(x_2)] (X_{22} + Y_{\max} Y_{22}), \quad (41)$$

where  $Y_{\max} = \tilde{m}_1/m_{\text{atm}}$ . One can see that if  $X_{22} = 1$  and  $Y_{22} = 0$ , corresponding to  $\Omega = R_{13}$ , then one recovers  $\xi_{\varepsilon_1} = 1$ , independently of the value of  $x_2$ . However, one can now perceive another possibility: if  $|X_{22} + Y_{\max} Y_{22}| \gg 1$ , then the  $CP$  asymmetry can be enhanced, i.e.  $\xi_{\varepsilon_1} > 1$ , even in the HL, when  $x_2 \gg 1$ . There are nevertheless some limitations. First of all, Yukawa couplings cannot be larger than  $\sim 0.1$  for the Eq. (16) to hold. Moreover, when calculating the final  $B - L$  asymmetry, one also has to take into account the increased wash-out. In the end, for fully hierarchical neutrinos, this possibility should be regarded as a very special case, also because it implies unnaturally huge phase cancellations due to the  $\Omega$  orthogonality. On the other hand, for quasi-degenerate light neutrinos this term can now be more easily dominant, since the first term is suppressed by  $\beta_{\max}(m_1, \tilde{m}_1) < 1$  [6, 13], while  $\Delta \varepsilon_1$  is not and thus an account of this term makes possible to evade the upper bound on the light neutrino masses [17].

## 5 Beyond the HL in strong-strong wash-out scenarios

In this section we discuss the effects arising in models beyond the HL and the conditions for the HL to be recovered. There is a large variety of possibilities and for definiteness we focus on a particularly interesting class that provides a useful framework to understand the general effects. We still assume a partial hierarchy, such that  $M_3 \gg M_1, M_2$ , while  $M_1$  and  $M_2$  can be arbitrarily close. This results in  $|\varepsilon_3| \ll |\varepsilon_1|, |\varepsilon_2|$  and the final asymmetry can be calculated as

$$N_{B-L}^f \simeq \varepsilon_1 \kappa_1^f + \varepsilon_2 \kappa_2^f. \quad (42)$$

Another convenient restriction is to focus on *strong-strong (ss) wash-out scenarios*, where both  $K_1$  and  $K_2 \gtrsim 5$ . In this way  $\kappa_1^f$  and  $\kappa_2^f$  can be calculated inserting the equilibrium values for both rate abundances into the general expressions (15), such that

$$\kappa_1^f(K_1, K_2, \delta_2) \simeq \kappa_1^{\text{ss}}(K_1, K_2, \delta_2) \equiv - \int_0^\infty dz' \frac{dN_{N_1}^{\text{eq}}}{dz'} e^{-\int_{z'}^\infty dz'' [W_1^{\text{ID}}(z'') + W_2^{\text{ID}}(z'')]} , \quad (43)$$



$$\kappa_2^f(K_1, K_2, \delta_2) \simeq \kappa_2^{ss}(K_1, K_2, \delta_2) \equiv - \int_0^\infty dz' \frac{dN_{N_2}^{\text{eq}}}{dz'} e^{-\int_{z'}^\infty dz'' [W_1^{\text{ID}}(z'') + W_2^{\text{ID}}(z'')]} . \quad (44)$$

In Fig. 5 we have plotted  $\kappa_1^{ss}(K_1, K_2, \delta_2)$  and  $\kappa_2^{ss}(K_1, K_2, \delta_2)$  for the indicated values of  $\delta_2$ , and for  $K_2 = K_{\text{atm}}$ . This value is particularly convenient to highlight the cumulative effect of the wash-out.

There are two simple limits where these integrals can be worked out explicitly. The first is the HL ( $\delta_2 \gg 1$ ). In this case the wash-out from  $N_2$  inverse decays acts only at  $z \ll z_B - 2$  and does not affect the dominant contribution produced around  $z \sim z_B$ . Therefore, one recovers the result valid in the  $N_1$ DS, i.e.  $\kappa_1^{ss}|_{\text{HL}} \simeq \kappa(K_1)$  (cf. (25)). It is also possible to obtain an analytic expression for  $\kappa_2^{ss}$  in the HL considering that the wash-out from  $N_1$  inverse decays does not interfere with the asymmetry production and with the wash-out from  $N_2$  decays and inverse decays respectively. Indeed, the wash-out from  $N_1$  inverse decays is active for  $z_{\text{on}}(K_1) \simeq 2/\sqrt{K_1} \lesssim z \lesssim z_{\text{off}}(K_1)$  and peaked at  $z_{\text{max}} \simeq 2.4$  [4], when the wash-out by  $N_2$  inverse decays is already off at  $z \simeq z_{\text{off}}(K_2)M_1/M_2 \ll z_{\text{on}}(K_1)$  and practically all  $N_2$ 's have already decayed. This results in a simple factorized analytic expression,

$$\kappa_2^{ss}|_{\text{HL}} \simeq \kappa(K_2) e^{-\int_0^\infty dz' W_1^{\text{ID}}(z')} \simeq \kappa(K_2) e^{-\frac{3\pi}{8} K_1} , \quad (45)$$

that is shown in Fig. 5 together with Eq. (25) for  $\kappa_1$  (circles). One can see how they well reproduce  $\kappa_1^{ss}$  and  $\kappa_2^{ss}$  in the HL. For  $\delta_2 > \delta_2^{\text{HL}}|_{\kappa_1^f}$  the HL for  $\kappa_1^f$  is recovered within 10%. In Fig. 6 we show  $\delta_2^{\text{HL}}|_{\kappa_1^f}$  as a function of  $K_1$  for fixed  $K_2 = K_{\text{atm}}$ . We also show (crosses) an analytic conservative estimate [8],

$$\delta_2 \geq \frac{z_B(K_2) + 2}{z_B(K_1) - 2} - 1 , \quad (46)$$

obtained neglecting the wash-out of the lightest RH neutrino on the asymmetry produced from the second lightest. One can see how indeed this is always more conservative, especially at large values of  $K_1$ , where the wash-out of the lightest RH is stronger. The second limit is the DL ( $M_2 = M_1$ , i.e.  $\delta_2 = 0$ ). This has been recently studied in [16] within the context of radiative leptogenesis. In this case one has  $z_2 = z_1 = z$ , so that  $dN_{N_1}^{\text{eq}}/dz' = dN_{N_2}^{\text{eq}}/dz'$  and  $W_1^{\text{ID}}/K_1 = W_2^{\text{ID}}/K_2$  (cf. (13)). It is then easy to derive a simple result,

$$\kappa_1^{ss}(K_1, K_2)|_{\delta_2=0} = \kappa_2^{ss}(K_1, K_2)|_{\delta_2=0} = \kappa(K_1 + K_2) , \quad (47)$$

indicating that in the DL the  $N_1$  and  $N_2$  wash-out contributions add up and tend to suppress the final asymmetry compared to the HL. This effect goes into the opposite direction compared to the enhancement of the  $CP$  asymmetry when masses get close. Therefore one has to be careful and check which effect is dominant between the two. As




Figure 5: Efficiency factors in the strong-strong wash-out case for  $K_2 = K_{\text{atm}}$ .




Figure 6: Plot of  $\delta_2^{\text{HL}}|_{\kappa_1^f}$  and of  $\delta_2^{\text{HL}}|_{N_{B-L}^f}$  vs.  $K_1$  for  $K_2 = K_{\text{atm}}$ . The crosses are the analytic estimate Eq. (46).

we will see a precise answer depends on the particular form of the orthogonal matrix and on the values of  $\delta_2$ ,  $K_1$  and  $K_2$ . One can see in Fig. 5 that the expression (47) reproduces well both  $\kappa_1^{\text{ss}}$  and  $\kappa_2^{\text{ss}}$  in the DL (open squares). Notice that this result is easily extended to the full DL ( $M_1 = M_2 = M_3$ ) with all  $K_i \gtrsim 5$ , where one obtains  $\kappa_i^f = \kappa(K_1 + K_2 + K_3)$ ,  $i = 1, 2, 3$ .

We have also worked out useful fits for any value of  $\delta_2$ ,

$$\kappa_1^{\text{fit}}(\delta_2) = \frac{2}{z_B(K_1 + K_2^{(1-\delta_2)^3})(K_1 + K_2^{1-\delta_2})} \quad (48)$$

and

$$\kappa_2^{\text{fit}}(\delta_2) = 2 \frac{\left[1 + 2 \ln\left(\frac{1+\delta_2}{1-\delta_2}\right)\right]^2}{z_B(K_2 + K_1^{(1-\delta_2)^3})(K_2 + K_1^{1-\delta_2})} \times e^{-\frac{3\pi}{8} K_1 \left(\frac{\delta_2}{1+\delta_2}\right)^{2.1}}. \quad (49)$$

In Fig. 5 one can see (black squares) that they fit well  $\kappa_1^{\text{ss}}$  and  $\kappa_2^{\text{ss}}$  for  $\delta_2 = 0.1$ . Notice moreover that the analytic expressions (25), (45) are correctly recovered in the HL and (47) in the DL.

We have thus seen that the simple class of strong-strong wash-out scenarios allows one to describe the most relevant features of models beyond the HL with simple analytic

expressions. More general cases can be easily understood and studied extending these results. In order to calculate the  $CP$  asymmetries and the lower bound on  $M_1$  in scenarios beyond the HL, we focus on two particularly interesting realizations of strong-strong wash-out scenarios: the first is  $\Omega = R_{13}$ , where the phase is maximal and an interesting result holds for the  $CP$  asymmetry  $\varepsilon_1$  beyond the HL, the second is the limit  $M_3 \gg 10^{14} \text{ GeV}$ , where the heaviest RH neutrino decouples and plays no role.

### 5.1 ‘Maximal phase’ scenario

Assuming  $\Omega = R_{13}$  (cf. (21)) and  $M_3 \gg M_1, M_2$  we have seen that the  $CP$  asymmetry  $\varepsilon_1$ , for hierarchical light neutrinos, is always given by the HL independently of  $\delta_2$ , i.e.  $\varepsilon_1 = \bar{\varepsilon}(M_1) \sin \delta_L^{(1)}$ , where the phase is maximal if  $X_{31} = 0$ . Moreover  $\varepsilon_2 = 0$ , so that the asymmetry is generated only from  $N_1$  decays. The only effect of  $N_2$  is the additional wash-out, described by a fixed value of the decay parameter  $\bar{K}_2 = K_{\text{sol}}$  for normal hierarchy and  $\bar{K}_2 = \sqrt{K_{\text{atm}}^2 - K_{\text{sol}}^2}$  for inverted hierarchy. The final asymmetry is then, for maximal phase, given by

$$N_{B-L}^f(M_1, K_1, \delta_2) = \bar{\varepsilon}(M_1) \kappa_1^f(K_1, \bar{K}_2, \delta_2). \quad (50)$$

For  $K_1 \gtrsim 5$  one can use for  $\kappa_1^f$  the analytic expressions obtained in the strong-strong wash-out regime and one obtains for the lower bounds,

$$M_1^{\min}(K_1, \delta_2) \simeq \frac{4.2 \times 10^8 \text{ GeV}}{\kappa_1^{\text{ss}}(K_1, \bar{K}_2, \delta_2)}, \quad (51)$$

and

$$T_{\text{in}}^{\min}(K_1, \delta_2) = \frac{M_1^{\min}(K_1, \delta_2)}{z_{\star}(K_1, \bar{K}_2, \delta_2)}, \quad (52)$$

where  $z_{\star}(K_1, K_2, \delta_2)$  is the  $T_{\text{in}}^{\min}$  relaxation factor and is approximately given, in the HL, by  $z_{\star}^{\text{HL}} \simeq z_B(K_1) - 2$  [4] and, in the DL, by  $z_{\star}^{\text{DL}} \simeq z_B(K_1 + K_2) - 2$ . For intermediate situations a good fit is given by  $z_{\star}(K_1, K_2, \delta_2) \simeq z_B(K_1 + K_2^{(1-\delta_2)^3}) - 2$ . In the top panel of Fig. 8 we compare the lower bounds valid in the HL with those for  $\delta_2 = 0.1$  (thick lines). One can see how the effect of the additional wash-out makes them more stringent. Moreover, one can see how the degeneracy between normal and inverted hierarchy in the HL is broken for finite value of  $\delta_2$  because of the different  $K_2$  value. In the bottom panel we have compared the lower bounds for  $\delta_2 = 0.1$  with those for  $\delta_2 = 0.01$  (thick lines). One can see that these sensibly change only at small  $K_1 \sim 10$ , while at larger values they are the same. This results from the fact that the effect of additional wash-out saturates and the DL is reached (cf. Fig. 5). Therefore, this model represents an interesting example of how going beyond the HL does not necessarily relaxes the lower bounds on  $M_1$  and  $T_{\text{in}}$ .

## 5.2 The limit of very heavy $N_3$

In the limit  $M_3 \gg 10^{14} \text{ GeV}$ , the orthogonal see-saw matrix  $\Omega$  necessarily reduces to a special form [21, 29, 30]

$$\Omega = \begin{pmatrix} 0 & 0 & 1 \\ \sqrt{1 - \Omega_{31}^2} & -\Omega_{31} & 0 \\ \Omega_{31} & \sqrt{1 - \Omega_{31}^2} & 0 \end{pmatrix}, \quad (53)$$

obtained from the general one for  $\Omega_{22}^2 = \Omega_{31}^2$  and  $\Omega_{22}^2 + \Omega_{31}^2 = 1$ . In terms of the parametrization with complex rotations this corresponds to  $w_{22} = 0$  and  $w_{21} = 1$ , implying  $\Omega_{21} = \sqrt{1 - w_{31}^2}$  (cf. (22)). This limit also implies  $m_1 \sim 0.01 \text{ eV} (10^{14} \text{ GeV}/M_3) (\text{Re}[U^\dagger h]_{13}^2/0.1) \ll m_{\text{sol}}$ , i.e. fully hierarchical light neutrinos. For normal hierarchy the values of  $K_1$  and  $K_2$  are given by

$$K_1 = K_{\text{sol}} \rho_{21} + K_{\text{atm}} \rho_{31} \quad \text{and} \quad K_2 = K_{\text{sol}} \rho_{31} + K_{\text{atm}} \rho_{21}, \quad (54)$$

so that  $K_1, K_2 \geq K_{\text{sol}} \simeq 9$  and  $K_1 + K_2 \geq K_{\text{sol}} + K_{\text{atm}} \simeq 60$ . For inverted hierarchy the same expressions hold with the replacement  $K_{\text{sol}} \rightarrow \sqrt{K_{\text{atm}}^2 - K_{\text{sol}}^2}$ . Therefore, in this scenario, both  $N_1$  and  $N_2$  decay in the strong wash-out regime, while the heaviest RH neutrino decouples completely. Notice moreover that  $\varepsilon_3 = 0$ , so that the final  $B - L$  asymmetry can be calculated as the sum of the two contributions from the two lightest RH neutrinos and it will depend on a set of 4 parameters that can be conveniently chosen to be  $M_1, K_1, K_2$  and  $\delta_2$ .

Let us calculate the two  $CP$  asymmetries  $\varepsilon_1$  and  $\varepsilon_2$ . As anticipated, it is easy to check that  $\Delta\varepsilon_1$ , defined in Eq. (27) for  $\varepsilon_1$ , vanishes. Therefore, the  $CP$  asymmetry enhancement is described just by the function  $\xi(x_2)$ , as in [13]. Moreover, since light neutrinos are fully hierarchical, one has  $\beta_{\text{max}}(m_1, \tilde{m}_1) = 1$  (cf. (30)). The expression (33) for the effective phase  $\sin \delta_L^{(1)}$  specializes to

$$\sin \delta_L^{(1)} = \frac{K_{\text{atm}}}{K_1} Y_{31} (1 - \sigma^2), \quad (55)$$

where  $Y_{31}$  has to be regarded as a function of  $K_1$  and  $K_2$ . Turning to  $\varepsilon_2$ , the general expression (16) can be re-cast as

$$\varepsilon_2 = \frac{3}{16\pi} \sum_{i=1,3} \frac{\text{Im}[(h^\dagger h)_{2i}^2] \xi(x_i/x_2)}{(h^\dagger h)_{22} \sqrt{x_i/x_2}}, \quad (56)$$

and using [8]

$$(h^\dagger h)_{ij} = \frac{\sqrt{M_i M_j}}{v^2} \sum_h m_h \Omega_{hi}^* \Omega_{hj}, \quad (57)$$

one can check that the term  $(h^\dagger h)_{23} = 0$ . After some algebraic manipulations, one finds

$$\varepsilon_2 \simeq \bar{\varepsilon}(M_2) \xi(1/x_2) \sin \delta_L^{(2)}. \quad (58)$$

We have introduced a second effective leptogenesis phase

$$\sin \delta_L^{(2)} \equiv -\frac{K_{\text{atm}}}{K_2} Y_{31} (1 - \sigma^2) = -\frac{K_1}{K_2} \sin \delta_L^{(1)}, \quad (59)$$

which always has opposite sign compared to  $\sin \delta_L^{(1)}$ . The final asymmetry (cf. (42)) can then be written as

$$N_{B-L}^f(M_1, K_1, K_2, \delta_2) = \bar{\varepsilon}(M_1) \sin \delta_L^{(1)}(K_1, K_2) \alpha(K_1, K_2, \delta_2), \quad (60)$$

where

$$\alpha(K_1, K_2, \delta_2) \equiv \xi(x_2) \kappa_1^{\text{ss}}(K_1, K_2, \delta_2) - \sqrt{x_2} \xi(1/x_2) \frac{K_1}{K_2} \kappa_2^{\text{ss}}(K_1, K_2, \delta_2). \quad (61)$$

Notice that in the HL one has

$$\frac{\varepsilon_2 \kappa_2^{\text{ss}}}{\varepsilon_1^{\text{HL}} \kappa_1^{\text{ss}}|_{\text{HL}}} \simeq -\sqrt{x_2} \xi(1/x_2) \frac{K_1}{K_2} \frac{\kappa_2^{\text{ss}}(K_1, K_2, \delta_2)}{\kappa(K_1)} \xrightarrow{\text{HL}} -\frac{2 \ln x_2}{3 \sqrt{x_2}} \frac{K_1^2}{K_2^2} \frac{z_B(K_1)}{z_B(K_2)} e^{-\frac{3\pi}{8} K_1} \ll 1. \quad (62)$$

As a consequence the contribution from the second lightest RH neutrino becomes negligible and one recovers the  $N_1$ DS. In Fig. 6 we show  $\delta_2^{\text{HL}}|_{N_{B-L}^f}$ , such that for  $\delta_2 > \delta_2^{\text{HL}}|_{N_{B-L}^f}$  the HL for the final asymmetry, in the case of very heavy  $N_3$ , is recovered within a 10% precision.

In Fig. 7 we give an example of the dynamics of the asymmetry generation with two RH neutrinos, a generalization of the example of Fig. 3 for the  $N_1$ DS. We show the quantity  $[d\kappa_1/dz' + (\varepsilon_2/\varepsilon_1) d\kappa_2/dz']_{z' \leq z}$  for  $K_1 = 10$  and  $K_2 = 50$ . In the top panel  $\sqrt{x_2} = M_2/M_1 = 10$  ( $\delta_2 = 9$ ). For this choice of the parameters, one has  $\varepsilon_2/\varepsilon_1 \simeq -0.035$ . At small values  $z \simeq 0.1$  the  $N_2$  decays give a non-negligible contribution to the total asymmetry but with negative sign. However, this contribution is completely washed-out at  $z \sim 1$  by the  $N_1$  inverse decays. Therefore, this example illustrates well how the HL is recovered for large  $\delta_2$ , where the production of the asymmetry and the wash-out from  $N_2$  and from  $N_1$  occur at two well-separated stages. This numerical example completes the qualitative discussion in [8] where the condition (46) was found. In the bottom panel we show an example for  $\delta_2 = 0.1$ , implying a minus sign in  $\xi(1/x_2)$  (cf. Fig. 1) that cancels with the minus sign in  $\sin \delta_L^{(2)}$ , such that now the asymmetry produced from the  $N_2$ 's has the same sign as the one produced from the  $N_1$ 's. Therefore, both the two productions and the two wash-out occur simultaneously and add up.





Figure 7: Dynamics of the asymmetry generation with 2 RH neutrinos.

Let us now calculate the lower bounds on  $M_1$  and on  $T_{\text{in}}$ . They can be written as

$$M_1^{\text{min}}(K_1, \delta_2) = \frac{4.2 \times 10^8 \text{ GeV}}{\sin \delta_L^{(1)}(K_1, K_2^*) \alpha(K_1, K_2^*, \delta_2)} \quad (63)$$

and

$$T_{\text{in}}^{\text{min}}(K_1, \delta_2) = \frac{M_1^{\text{min}}(K_1, \delta_2)}{z_*(K_1, K_2^*, \delta_2)}, \quad (64)$$

where  $K_2^*$  is the value of  $K_2$  that maximizes  $N_{B-L}^f$  (cf. (60)). In the HL one has  $\alpha(K_1, K_2^*, \delta_2 \gg 1) = \kappa_1^{\text{ss}}|_{\text{HL}} \simeq \kappa(K_1)$ , such that the  $N_1$ DS is recovered but the effective phase is not maximal (cf. (55)) and the lower bounds are more stringent [30]. Maximizing the effective phase  $\sin \delta_L^{(1)}(K_1, K_2)$  over  $K_2$ , one obtains  $\sin \delta_L^{(1)} = f(m_2, \tilde{m}_1) (1 - \sigma)$  [8]. In the case of normal hierarchy  $\sigma \simeq m_{\text{sol}}/m_{\text{atm}} \simeq 0.19$ . The function  $f(m_{\text{sol}}, \tilde{m}_1)$  can then be calculated using Eq. (32) with the replacement  $m_1 \rightarrow m_{\text{sol}} \ll m_{\text{atm}}$ . In this way, from Eq. (36) for  $(M_1^{\text{min}})_{\text{HL}}$ , one obtains

$$(M_1^{\text{min}})_{\text{HL}} \simeq \frac{4.2 \times 10^8 \text{ GeV}}{\kappa(K_1) f(m_{\text{sol}}, \tilde{m}_1) (1 - \sigma)}. \quad (65)$$

This lower bound and the associated one on  $T_{\text{in}}^{\text{min}}$  are shown in Fig. 4 (thin lines) and compared with those obtained for  $\sin \delta_L^{(1)} = 1$  (thick lines). For inverted hierarchy one has  $\sigma \simeq \sqrt{1 - (m_{\text{sol}}/m_{\text{atm}})^2}$  and  $f(m_2, \tilde{m}_1) \simeq \sqrt{1 - (m_2/\tilde{m}_1)^2}$ . The bounds become much more restrictive [30] for two reasons: first now  $K_1 \gtrsim K_{\text{atm}}$ , second there is strong phase cancellation in  $\sin \delta_L^{(1)}$ . This actually occurs on more general grounds [8] and one can say that for inverted hierarchy the constraints are more stringent than for normal hierarchy, except for the two special cases already mentioned in Section 3. The first one is for  $\Omega = R_{13} R_{23}$  ( $\Omega_{21} = 0$ ) and  $Y_{31} = Y_{\text{max}}$ , where the phase is maximal and one obtains the most conservative lower bounds on  $M_1$  and  $T_{\text{in}}$  in the HL, the same for normal and inverted hierarchy. As soon as one diverges from this special case, the degeneracy gets broken. A second special case is for  $\Omega = R_{12} R_{23}$  ( $\Omega_{31} = 0$ ) and  $Y_{21} = Y_{\text{max}}/\sigma$ , so that  $\sin \delta_L^{(1)} = \sigma$ , larger for inverted hierarchy than for the normal one. However, as we said, the wash-out is stronger for inverted hierarchy than for normal hierarchy and the two things compensate, such that the lower bounds are practically equivalent, though not equal. Except for these two cases the allowed region in the space parameters is larger for normal hierarchy than for inverted hierarchy and leptogenesis tends to favor normal over inverted one. The case of very large  $M_3$  represents the most extreme one in this respect. Also in the  $N_2$ DS the lower bounds on  $M_2$  and  $T_{\text{in}}$  are more stringent in the case of inverted hierarchy than in normal hierarchy. We can thus conclude that if the light



neutrino hierarchy were to be inverted, then the allowed region would be considerably reduced.

Let us now see what happens beyond the HL, i.e. for small values of  $\delta_2$ . There are three effects, all accounted for by the function  $\alpha(K_1, K_2, \delta_2)$  (cf. (61)). The first is that the asymmetry produced from the  $N_2$ 's becomes non-negligible and since  $\xi(1/x_2)$  becomes negative at small  $\delta_2$ , cancelling the minus sign in  $\sin \delta_L^{(2)}$ , the two asymmetries add up. However, this effect brings a factor 2 enhancement at most [13], while the major effect is the enhancement of both  $\varepsilon_1$  and  $\varepsilon_2$ , described by  $\xi(x_2)$  for  $\varepsilon_1$  and by  $\xi(1/x_2)$  for  $\varepsilon_2$ . The third effect is that also the two wash-out effects add up. As we have seen (cf. Fig. 5) this effect saturates at  $\delta_2 \simeq 0.01$  and goes into the opposite direction compared to the other two, thus contrasting the enhancement of the asymmetry for moderately small values  $\delta_2 \sim 0.1$ . This is clearly visible in the top panel of Fig. 8, where we have compared the results holding in the HL with those obtained for  $\delta_2 = 0.1$ . The lower bounds for the very heavy  $N_3$  case are indicated with thin lines. One can see how these are relaxed just by a factor 2 instead of one order of magnitude as one could expect from  $CP$  asymmetry enhancement plus, sub-dominantly, the contribution to the final asymmetry from  $N_2$  decays. However, if one considers the DL, then the asymmetry and the wash-out cumulative effects saturate, while the  $CP$  asymmetry enhancement lowers the bounds  $\propto \xi(x_2) \simeq (3\delta_2)^{-1}$  and this can be clearly described analytically. Indeed, as it can be seen from Fig. 1, for small  $\delta_2$  one has  $\xi(1/x_2) = -\xi(x_2)$  and therefore, in the DL, the expression for the final asymmetry simplifies to (cf. (60), (55) and (25))

$$(N_{B-L}^f)_{\text{DL}}(M_1, K_1, K_2) \equiv 2(1 - \sigma^2) K_{\text{atm}} \bar{\varepsilon}(M_1) \xi(x_2) z_B(K_1 + K_2) \frac{Y_{31}(K_1, K_2)}{K_1 K_2} \propto \xi(x_2). \quad (66)$$

In the bottom panel of Fig. 8 we compared the lower bounds for  $\delta_2 = 0.1$  with those for  $\delta_2 = 0.01$  and one can see how the lower bound on  $M_1$  ( $T_{\text{in}}$ ) is  $4 \times 10^{10}$  GeV ( $5 \times 10^9$  GeV) for  $\delta_2 = 0.1$  and lowers to  $4 \times 10^9$  GeV ( $5 \times 10^8$  GeV) for  $\delta_2 = 0.01$ . For  $\delta_2 \lesssim 0.01$ , the wash-out and asymmetry cumulative effects are negligible and the  $CP$  asymmetry enhancement is the only left one, so that

$$(M_1^{\text{min}})_{\text{DL}} \simeq 4 \times 10^9 \text{ GeV} \left( \frac{0.01}{\delta_2} \right) \quad \text{and} \quad (T_{\text{in}}^{\text{min}})_{\text{DL}} \simeq 5 \times 10^8 \text{ GeV} \left( \frac{0.01}{\delta_2} \right). \quad (67)$$

The enhancement at small  $\delta_2$  has an upper limit set by the condition of validity of the Eq. (16), which implies that values  $|\varepsilon_1| \sim \mathcal{O}(0.01)$  can always be reached and thus that there is no lower bound on  $M_1$  if arbitrarily small  $\delta_2$  values are possible. Examples of models where such small values can be motivated are soft leptogenesis [31] and radiative leptogenesis [16].




Figure 8: Lower bounds on  $M_1$  and  $T_{in}$  in the case of ‘maximal phase’ (thick lines) and very large  $M_3$  (thin lines). Upper panels: comparison between the HL ( $\delta_2 \gg \delta_2^{\text{HL}}$ ) and  $\delta_2 = 0.1$ . Lower panels: comparison between  $\delta_2 = 0.1$  and  $\delta_2 = 0.01$ .

In conclusion a strong relaxation of the lower bounds on  $M_1$  and  $T_{\text{in}}$  is attainable only for degeneracies  $\delta_2 \ll 0.1$ . This confirms and actually strengthens the conclusions of [13], showing at the same time that effects arising from a term  $\Delta\varepsilon_1$  are possible only for a very special category of models where a complex rotation  $R_{23}$  is switched on with very high values of  $|\Omega_{22}|$  implying unlikely phase cancellations. This term can nevertheless play an interesting role in relaxing the upper bound on the absolute neutrino mass scale [17].

## 6 Final Discussion

We have calculated the final baryon asymmetry in leptogenesis scenarios beyond the HL. Our main result is that the lower bounds on  $M_1$  and  $T_{\text{in}}$  can be relaxed but not necessarily, depending on the particular form of the see-saw orthogonal matrix and on the value of  $\delta_2$ . Very small values  $\delta_2 \ll 0.1$  are necessary to have a significant relaxation of the lower bounds, confirming and strengthening what was found in [13]. One should then understand, on theoretical grounds, whether such small values of  $\delta_2$  should be considered natural or not.

There is no compelling reason to exclude  $M_1$  as large as  $10^{11}$  GeV and  $T_{\text{in}}$  as large as  $10^{10}$  GeV. In many specific models the lower bound on  $M_1$  is however a problem, especially in those cases where the Dirac neutrino mass matrix is required to resemble the up quark mass matrix [32]. In supergravity the gravitino problem requires reheating temperatures that can be quite difficultly compatible with the lower bound from leptogenesis in the HL. In this case  $\delta_2 \sim 10^{-3}$  or smaller would be desirable. Within a minimal model with 3 RH neutrinos, the  $N_2$ DS is an alternative solution. Here one does not need to go beyond the HL for  $M_1$  to be arbitrarily low, but  $T_{\text{in}} \gtrsim 5 \times 10^9$  GeV still applies. At the moment there is no reason to prefer one of the two possible ways to go beyond the  $N_1$ DS, in any case still the most appealing scenario. As we have seen, a  $N_2$ DS is not possible for very large  $M_3$ , since in this limit and for large  $\delta_2 \gg 1$  the  $CP$  asymmetry  $\varepsilon_2$  vanishes. It is important to notice moreover that the  $N_2$ DS will be disproved if the absolute neutrino mass scale is found to be  $m_1 \gtrsim (10^{-3} - 10^{-2})$  eV. On the other hand, we do not have currently any experimental test to disprove small  $\delta_2$  values and there is a large freedom in the choice of the heavy neutrino mass spectrum, also on pure theoretical grounds.

It is quite interesting that a normal hierarchical light neutrino spectrum is desirable since an inverted one exacerbates the problem with the  $M_1$  and  $T_{\text{in}}$  lower bounds.

Notice that flavor effects [18, 19, 20] introduce new interesting modifications of the lower bounds but they do not change any of the conclusions we have discussed here and will be addressed in a forthcoming paper.

A test of leptogenesis in the next years represents a great challenge. However, in this respect, it is more convenient to regard it within the wider context of a test of the see-saw mechanism, where leptogenesis is one among various phenomenological consequences.

A smoking gun seems possible only in the very fortunate case that the lightest or the two lightest heavy neutrinos are at the TeV scale and possess extra gauge interactions, making possible their production and detection at the LHC. Because the lower bound on  $M_1$  is so large, this cannot happen in the HL within the usual  $N_1$ DS but it becomes possible either in the DL with  $\delta_2 \lesssim 10^{-7}$  [33] or, more appealingly, in the  $N_2$ DS [8]. We can thus conclude that exciting results are likely to come again in leptogenesis in the near future.

### Acknowledgments

It is a pleasure to thank A. Anisimov, A. Broncano, G. Branco, R. Gonzalez Felipe, S. Petcov, M. Plümacher, G. Raffelt and A. Weiler for useful discussions.

## References

- [1] M. Fukugita, T. Yanagida, Phys. Lett. **B 174** (1986) 45 .
- [2] P. Minkowski, Phys. Lett. B **67** (1977) 421; T. Yanagida, in *Workshop on Unified Theories*, KEK report 79-18 (1979) p. 95; M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity* (North Holland, Amsterdam, 1979) eds. P. van Nieuwenhuizen, D. Freedman, p. 315; S.L. Glashow, in *1979 Cargese Summer Institute on Quarks and Leptons* (Plenum Press, New York, 1980) p. 687; R. Barbieri, D. V. Nanopoulos, G. Morchio and F. Strocchi, Phys. Lett. B **90** (1980) 91; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912.
- [3] E. W. Kolb and M. S. Turner, Ann. Rev. Nucl. Part. Sci. **33** (1983) 645.
- [4] W. Buchmüller, P. Di Bari and M. Plümacher, Annals Phys. **315** (2005) 305.
- [5] T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, Phys. Lett. B **464** (1999) 12.
- [6] S. Davidson, A. Ibarra, Phys. Lett. **B 535** (2002) 25.
- [7] W. Buchmüller, P. Di Bari and M. Plümacher, Nucl. Phys. B **643** (2002) 367.
- [8] P. Di Bari, Nucl. Phys. B **727** (2005) 318.

- [9] G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, Nucl. Phys. B **685** (2004) 89.
- [10] P. Di Bari, arXiv:hep-ph/0406115.
- [11] G. F. Giudice, E. W. Kolb and A. Riotto, Phys. Rev. D **64** (2001) 023508.
- [12] W. Buchmüller, P. Di Bari and M. Plümacher, Phys. Lett. B **547** (2002) 128.
- [13] W. Buchmüller, P. Di Bari and M. Plümacher, Nucl. Phys. B **665** (2003) 445.
- [14] M. Plümacher, Z. Phys. C **74** (1997) 549.
- [15] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim, I. Masina, M. N. Rebelo and C. A. Savoy, Phys. Rev. D **67** (2003) 073025; A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. B **692** (2004) 303; T. Hambye, Y. Lin, A. Notari, M. Papucci and A. Strumia, Nucl. Phys. B **695** (2004) 169.
- [16] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim and B. M. Nobre, Phys. Lett. B **633** (2006) 336.
- [17] T. Hambye, Y. Lin, A. Notari, M. Papucci and A. Strumia, Nucl. Phys. B **695** (2004) 169.
- [18] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, Nucl. Phys. B **575** (2000) 61.
- [19] T. Endoh, T. Morozumi and Z. h. Xiong, Prog. Theor. Phys. **111** (2004) 123; E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601** (2006) 164.
- [20] A. Abada, S. Davidson, F. X. Josse-Michaux, M. Losada and A. Riotto, arXiv:hep-ph/0601083.
- [21] P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548** (2002) 119.
- [22] M. A. Luty, Phys. Rev. D **45** (1992) 455.
- [23] A.D. Dolgov, Sov. J. Nucl. Phys. **32**, 831 (1980); E. W. Kolb, S. Wolfram, Nucl. Phys. B **172** (1980) 224.
- [24] W. Buchmüller, M. Plümacher, Phys. Lett. B **431** (1998) 354; A. Anisimov, A. Broncano and M. Plümacher, Nucl. Phys. B **737** (2006) 176.

- [25] M. Flanz, E. A. Paschos, U. Sarkar, Phys. Lett. **B 345** (1995) 248; Phys. Lett. **B 384** (1996) 487 (E) L. Covi, E. Roulet, F. Vissani, Phys. Lett. **B 384** (1996) 169
- [26] J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001) 171.
- [27] WMAP Collaboration, D. N. Spergel *et al.*, Astrophys. J. Suppl. **148** (2003) 175; M. Tegmark *et al.* [SDSS Collaboration], Phys. Rev. D **69** (2004) 103501.
- [28] O. Vives, arXiv:hep-ph/0512160.
- [29] A. Ibarra and G. G. Ross, Phys. Lett. B **575** (2003) 279.
- [30] P. H. Chankowski and K. Turzyski, Phys. Lett. B **570** (2003) 198.
- [31] G. D'Ambrosio, G. F. Giudice and M. Raidal, Phys. Lett. B **575** (2003) 75.
- [32] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim and M. N. Rebelo, Nucl. Phys. B **640** (2002) 202; S. F. King, Rept. Prog. Phys. **67** (2004) 107; E. K. Akhmedov, M. Frigerio and A. Y. Smirnov, JHEP **0309** (2003) 021.
- [33] A. Pilaftsis, Phys. Rev. D **56** (1997) 5431; S. F. King and T. Yanagida, Prog. Theor. Phys. **114** (2006) 1035.